

hep-th/9906023

YITP-99-31

May, 1999

A - D - E Singularity and the Seiberg-Witten Theory*

Katsushi Ito

*Yukawa Institute for Theoretical Physics
Kyoto University, Kyoto 606-8502, Japan*

Abstract

We study the low-energy effective theory of $N = 2$ supersymmetric Yang-Mills theory with ADE gauge groups in view of the spectral curves of the periodic Toda lattice and the A - D - E singularity theory. We examine the exact solutions by using the Picard-Fuchs equations for the period integrals of the Seiberg-Witten differential. In particular, we find that the superconformal fixed point in the strong coupling region of the Coulomb branch is characterized by the ADE superpotential. We compute the scaling exponents, which agree with the previous results.

* Talk given at the workshop “Gauge Theory and Integrable Models”, YITP, Kyoto, January 26–29, 1999

1 Introduction

Seiberg and Witten showed that the low-energy effective theory of $N = 2$ supersymmetric gauge theory in four dimensions is determined by the prepotential, a holomorphic function of the period integrals of the meromorphic one-form (the Seiberg-Witten differential) on a Riemann surface[1]. For a simple Lie group G , it has been proposed in [2, 3] that the spectral curve of the periodic Toda lattice associated with the dual affine Lie algebra $(\hat{G})^\vee$ provides the Riemann surface which describes the Coulomb branch of $N = 2$ supersymmetric Yang-Mills theory with the gauge group G . In the case of gauge theories with some matter hypermultiplets, the spectral curves and related integrable systems are discussed in [4]. Other systematic approaches based on the heterotic/type II duality[5] or the M5 branes [6] are also studied extensively.

In the present work we will study the exact solution of the low-energy effective theory from the viewpoint of the spectral curve of the periodic Toda lattice. For ADE type gauge groups, the spectral curves are shown to be the sum of the superpotential of two-dimensional topological Landau-Ginzburg models of ADE type and that of the topological CP^1 model. In a series of papers[7, 8, 9, 10], we have studied various aspects of the exact solution of the Seiberg-Witten theories with ADE gauge groups by using two-dimensional topological field theories.

This paper is organized as follows: In sect. 2, we introduce the spectral curve of the periodic Toda lattice associated with the dual of the affine Lie algebra $(\hat{G})^\vee$ for the gauge theory with gauge group G . In sect. 3, we consider the ADE gauge groups and express the spectral curve as the sum of the superpotential of the topological CP^1 model and the ADE minimal model. Using the flat coordinates in the A - D - E singularity theory, we derive the Picard-Fuchs differential equations obeyed by the period integral of the Seiberg-Witten differential. We then show that these equations are equivalent to the Gauss-Manin system for the ADE minimal model and the CP^1 model and the scaling relation for the Seiberg-Witten differential. In sect. 4, we study an exact solution in the strong coupling region. Argyres and Douglas[11] showed that there exists a non-trivial RG fixed point in the Coulomb branch such that the massless solitons with mutually non-local charges coexist and the theory corresponds to $N = 2$ superconformal field theory,

where the gauge invariant order parameters have fractional dimensions with respect to the BPS mass. We investigate this Argyres-Douglas point in the Coulomb branch of the $N = 2$ supersymmetric Yang-Mills theory for ADE gauge groups.

2 Spectral Curves and A - D - E Singularity

The low-energy properties of the Coulomb branch of $N = 2$ supersymmetric gauge theories with gauge group G are exactly described by holomorphic data associated with certain algebraic curves. In particular, the BPS mass formula is expressed in terms of the period integrals of the so-called Seiberg-Witten (SW) differential λ_{SW} :

$$m_{BPS} = |n^I a_I + m^I a_{DI}| \quad (1)$$

where n^I and m^I are integers and

$$a_I = \oint_{A_I} \lambda_{SW}, \quad a_{DI} = \oint_{B_I} \lambda_{SW}, \quad I = 1, \dots, r \quad (2)$$

along one-cycles A_I , B_I with appropriate intersections on the curve. Here r is the rank of the gauge group G . The low-energy effective theory effective action is described by the prepotential $\mathcal{F}(a)$. The dual period a_{DI} is then given by $\partial\mathcal{F}(a)/\partial a_I$.

We define the spectral curve for the periodic Toda lattice for the (twisted) affine Lie algebra $(\widehat{G})^\vee$. Let G be a simple Lie algebra with rank r . Let $\alpha_1, \dots, \alpha_r$ be simple roots of the Lie algebra and $\alpha_0 = -\theta$, where θ denotes the highest root. We consider a representation $\mathcal{R}$ with d dimensions. Let E_α be generators associated with the roots α and H^i those of the Cartan subalgebra, which are realized by $d \times d$ matrices in the representation $\mathcal{R}$. Introduce matrices A and B by

$$\begin{aligned} A(z) &= \sum_{i=1}^r b_i H_i + a_i (E_{\alpha_i} + E_{-\alpha_i}) + a_0 (z E_{\alpha_0} + z^{-1} E_{-\alpha_0}) \\ B(z) &= \sum_{i=1}^r b_i H_i + a_i (E_{\alpha_i} - E_{-\alpha_i}) + a_0 (z E_{\alpha_0} - z^{-1} E_{-\alpha_0}), \end{aligned} \quad (3)$$

where z is called as the spectral parameter. The equation of motion of the periodic Toda lattice is defined in the Lax form

$$\frac{dA}{dt} = [A, B]. \quad (4)$$

The spectral curve is defined by the characteristic polynomial of the matrix $A(z)$:

$$\mathcal{P}_G^{\mathcal{R}}(x; u_1, \dots, u_r, z) \equiv \det_{\mathcal{R}}(x1_d - A(z)) = 0. \quad (5)$$

Here u_i ($i = 1, \dots, r$) are the i -th order Casimirs of G with degree q_i where q_i is order of the Casimirs. The exponents e_i of G is related to q_i by $q_i = e_i + 1$. The second order Casimir u_2 has degree 2 and the top Casimir u_r has degree $q_r = h$, where h is the Coxeter number. The curve depends on the scale parameter $\mu^2 = \prod_{i=0}^r a_i^{n_i}$, where non-negative integers n_i 's are the Dynkin labels of the affine roots. The spectral parameter z and μ have degree $h^\vee$, the dual Coxeter number of G . The exponents and the (dual) Coxeter number are given in the table 1. The spectral curve is invariant under the transformation

group G	$(\tilde{G})^\vee$	h	$h^\vee$	exponents
A_r	$A^{(1)}$	$r + 1$	$r + 1$	$1, 2, \dots, r$
B_r	$A_{2r-1}^{(2)}$	$2r$	$2r - 1$	$1, 3, \dots, 2r - 1$
C_r	$D_{r+1}^{(2)}$	$2r$	$r + 1$	$1, 3, \dots, 2r - 1$
D_r	$D_r^{(1)}$	$2r - 2$	$2r - 2$	$1, 3, \dots, 2r - 3, r - 1$
E_6	$E_6^{(1)}$	12	12	1, 4, 5, 7, 8, 11
E_7	$E_7^{(1)}$	18	18	1, 5, 7, 9, 11, 13, 17
E_8	$E_8^{(1)}$	30	30	1, 7, 11, 13, 17, 19, 23, 29
F_4	$E_6^{(2)}$	12	9	1, 5, 7, 11
G_2	$D_4^{(3)}$	6	4	1, 5

Table 1: List of the (dual) Coxeter numbers and exponents

$z \rightarrow \mu^2/z$. For simply laced Lie algebra G , we have $h = h^\vee$. Therefore the top Casimir u_r and the spectral parameter z or its dual μ^2/z have the same degree. It is found that for the representation $\mathcal{R}$ of $G = ADE$, the spectral curve is given by

$$P_G^{\mathcal{R}}(x; u_1, \dots, u_r + z + \frac{\mu^2}{z}) = 0, \quad (6)$$

where $P_G^{\mathcal{R}}(x; u_1, \dots, u_r)$ is the characteristic polynomial of the representation $\mathcal{R}$ of G and expressed in the form of

$$P_G^{\mathcal{R}}(x; u_1, \dots, u_r) = \prod_{i=1}^d (x - \lambda_i \cdot a). \quad (7)$$

Here λ_i denote the weight vector of the representation $\mathcal{R}$. We list the explicit form of the spectral curve for A_r , D_r and E_6 Lie algebras for the d -dimensional representation $\underline{d}$:

- $A_r (A_r^{(1)}, \underline{r+1})$

$$x^{r+1} - u_1 x^{r-1} - \dots - u_r - \left(z + \frac{\mu^2}{z} \right) = 0 \quad (8)$$

- $D_r (D_r^{(1)}, \underline{2r})$

$$x^{2r} - u_1 x^{2r-2} - \dots - u_{r-2} x^4 - u_r x^2 - u_{r-1}^2 - x^2 \left(z + \frac{\mu^2}{z} \right) = 0 \quad (9)$$

- $E_6 (E_6^{(1)}, \underline{27})$ [12]

$$\frac{1}{2} x^3 \left(z + \frac{\mu^2}{z} + u_6 \right)^2 - q_1(x) \left(z + \frac{\mu^2}{z} + u_6 \right) + q_2(x) = 0, \quad (10)$$

where

$$\begin{aligned} q_1 &= 270x^{15} + 342u_1x^{13} + 162u_1^2x^{11} - 252u_2x^{10} + (26u_1^3 + 18u_3)x^9 \\ &\quad - 162u_1u_2x^8 + (6u_1u_3 - 27u_4)x^7 - (30u_1^2u_2 - 36u_5)x^6 + (27u_2^2 - 9u_1u_4)x^5 \\ &\quad - (3u_2u_3 - 6u_1u_5)x^4 - 3u_1u_2^2x^3 - 3u_2u_5x - u_2^3, \\ q_2 &= \frac{1}{2x^3}(q_1^2 - p_1^2p_2), \\ p_1 &= 78x^{10} + 60u_1x^8 + 14u_1^2x^6 - 33u_2x^5 + 2u_3x^4 - 5u_1u_2x^3 - u_4x^2 - u_5x - u_2^2, \\ p_2 &= 12x^{10} + 12u_1x^8 + 4u_1^2x^6 - 12u_2x^5 + u_3x^4 - 4u_1u_2x^3 - 2u_4x^2 + 4u_5x + u_2^2. \end{aligned} \quad (11)$$

For simply-laced Lie algebra G , $\widehat{G}$ is self-dual, i.e. $(G^{(1)})^\vee = G^{(1)}$. For non-simply laced Lie algebras, we have $\widehat{B}_r^\vee = A_{2r-1}^{(2)}$, $\widehat{C}_r^\vee = D_{r+1}^{(2)}$, $\widehat{F}_4^\vee = E_6^{(2)}$ and $\widehat{G}_2^\vee = D_4^{(3)}$. Thus we need the twisted affine Lie algebra to construct the spectral curve. The characteristic polynomial can be obtained by folding procedure of the corresponding Dynkin diagram. Due to $h \neq h^\vee$, the spectral parameter z or its dual μ^2/z appears in the spectral curve in a nontrivial way. Now we will write the explicit form of the spectral curves for d -dimensional representation $\underline{d}$ of non-simply laced Lie algebra G .

- For B_r ($A_{2r-1}^{(2)}, \underline{2r}$) case, the spectral curve of the representation $\underline{2r}$ is obtained from the characteristic polynomial $P_{A_{2r-1}}^{2r}(x; u_1, \dots, u_{2r-1})$ by the restriction $u_2 = \dots = u_{2r-4} = 0$ and $u_{2r-2} = z + \mu^2/z$:

$$x^{2r} - u_1 x^{2r-2} - \dots - u_{2r-1} - x \left(z + \frac{\mu^2}{z} \right) = 0. \quad (12)$$

- For C_r ($D_{r+1}^{(2)}; \underline{2r+2}$) case, the curve is obtained from the characteristic polynomial $P_{D_{r+1}}^{2r+2}(x; u_1, \dots, u_r, u_{r+1})$ by restricting $u_r = x - \mu^2/z$;

$$x^{2r+2} - u_1 x^{2r} - \dots - u_{r-1} x^4 - u_{r+1} x^2 - \left(z - \frac{\mu^2}{z} \right)^2 = 0. \quad (13)$$

- For F_4 ($E_6^{(2)}; \underline{27}$) case, the curve is obtained from the characteristic polynomial $P_{E_6}^{27}(x; u_1, u_2, u_3, u_4, u_5, u_6)$ by the restriction $u_2 = 0$ and $u_5 = -6(z + \mu^2/z)$;

$$\begin{aligned} & -8 \left(z + \frac{\mu^2}{z} \right)^3 + a_1(x) \left(z + \frac{\mu^2}{z} \right)^2 + a_2(x) \left(z + \frac{\mu^2}{z} \right) + a_3(x) = 0, \\ a_1(x) &= -636x^9 - 300u_1x^7 - 48u_1^2x^5 - 5u_3x^3 + 2u_4x, \\ a_2(x) &= -168x^{18} - 348u_1x^{16} - 276u_1^2x^{14} + (-116u_1^3 + 14u_3)x^{12} \\ & \quad + (-92u_4 - 20u_1^4 - 8u_1u_3)x^{10} + (-42u_1u_4 - 6u_1^2u_3)x^8 \\ & \quad + (-4u_6 - \frac{10}{3}u_1^2u_4 - \frac{2}{3}u_3^2)x^6 + (\frac{1}{3}u_3u_4 - \frac{2}{3}u_6u_1)x^4, \\ a_3(x) &= x^{27} + 6u_1x^{25} + 15u_1^2x^{23} + (20u_1^3 + u_3)x^{21} + (5u_4 + 4u_1u_3 + 15u_1^4)x^{19} \\ & \quad + (6u_1^2u_3 + 12u_1u_4 + 6u_1^5)x^{17} + (\frac{1}{3}u_3^2 + 5u_6 + 4u_1^3u_3 + \frac{26}{3}u_1^2u_4 + u_1^6)x^{15} \\ & \quad + (\frac{4}{3}u_1^3u_4 + \frac{19}{3}u_6u_1 + u_1^4u_3 + \frac{4}{3}u_3u_4 + \frac{2}{3}u_3^2u_1)x^{13} \\ & \quad + (\frac{1}{3}u_1^2u_3^2 - \frac{1}{3}u_1^4u_4 - \frac{15}{4}u_4^2 + 3u_6u_1^2)x^{11} \\ & \quad + (\frac{1}{3}u_6u_3 - \frac{4}{9}u_1^2u_3u_4 + \frac{1}{27}u_3^3 - \frac{13}{6}u_4^2u_1 + \frac{13}{27}u_6u_1^3)x^9 \\ & \quad + (-\frac{1}{9}u_3^2u_4 - \frac{1}{2}u_6u_4 + \frac{1}{9}u_6u_1u_3 - \frac{7}{36}u_1^2u_4^2)x^7 + (\frac{1}{12}u_4^2u_3 - \frac{1}{6}u_6u_1u_4)x^5 \\ & \quad + (-\frac{1}{54}u_4^3 - \frac{1}{108}u_6^2)x^3. \end{aligned} \quad (14)$$

- For G_2 ($D_4^{(3)}; \underline{8}$) case[3], the spectral curve is obtained from $P_{D_4}^8(x; u_1, u_2, u_3, u_4)$ by the restriction $u_1 = 2u$, $u_2 = -u^2 - z + \mu^2/z$, $u_3 = \sqrt{3}(z - \mu^2/z)$ and $u_4 =$

$$v + 2u(z + \mu^2/z);$$

$$3 \left(z - \frac{\mu^2}{z} \right)^2 - x^8 + 2ux^6 - \left[u^2 + \left(z + \frac{\mu^2}{z} \right) \right] x^4 + \left[v + 2u \left(z + \frac{\mu^2}{z} \right) \right] x^2 = 0. \quad (15)$$

We may write the spectral curve in the form of

$$z + \frac{\mu^2}{z} = W_G^{\mathcal{R}}(x, u_1, \dots, u_r), \quad (16)$$

namely we regard the curve as an fibration over CP^1 with the fiber characterized by the function $W_G^{\mathcal{R}}(x)$. For example, in the case of A_r , D_r and E_6 gauge groups, we obtain

- A_r

$$W_{A_r}^{r+1} = x^{r+1} - u_1 x^r - \dots - u_{r-1} x - u_r, \quad (17)$$

- D_r

$$W_{D_r}^{2r} = x^{2r-2} - u_1 x^{2r-4} - \dots - u_{r-2} x^2 - \frac{u_{r-1}^2}{x^2} - u_r, \quad (18)$$

- E_6

$$W_{E_6}^{27} = \frac{1}{x^3} (q_1 \pm p_1 \sqrt{p_2}) - u_6, \quad (19)$$

where p_1 , p_2 and q_1 are given in (11).

It is important to notice that $W_{ADE}(x)$ is nothing but the superpotential of the two-dimensional topological Landau-Ginzburg (LG) model of type ADE . $W_{A_r}^{r+1}(x)$ and $W_{D_r}^{2r}(x)$ are familiar superpotentials for the A_r and D_r type minimal models. For E_6 case, the function $W_{E_6}^{27}$ looks like very different from the usual deformation of the E_6 singularity written in terms of three variables

$$W_{E_6}(x_1, x_2, x_3) = x_1^4 + x_2^3 + x_3^2. \quad (20)$$

It is, however, found in [13] that $W_{E_6}^{27}$ is a single-variable version of the LG superpotential for the E_6 minimal model. On the other hand, the singularity of the form of (20) is obtained by considering the fibration of ALE spaces[5]. The relation of these two description of the Seiberg-Witten curves are discussed in [12].

For non-simply laced cases, we have rather nontrivial $W_G(x)$ from the spectral curves although in the two dimensional case, the LG superpotentials are obtained from the simply laced one by the folding procedure[14]. The functions $W_G^{\mathcal{R}}(x)$ are given as follows:

- B_r

$$W_{B_r}^{2r}(x; u_1, \dots, u_r) = \frac{W_{BC}(x; u_1, \dots, u_r)}{x}, \quad (21)$$

where the LG potential of BC type

$$W_{BC}(x; u_1, \dots, u_r) = x^{2r} - \sum_{i=1}^r u_i x^{2r-2i} \quad (22)$$

is obtained from the A_{2r-1} superpotential $W_{A_{2r-1}}(x; \tilde{u}_1, \dots, \tilde{u}_{2r-1})$ by the restriction $\tilde{u}_{2k} = 0$ ($k = 1, \dots, r-1$) and setting $u_k = \tilde{u}_{2k-1}$ ($k = 1, \dots, r$).

- C_r

$$W_{C_r}^{2r+2}(x; u_1, \dots, u_r) = \left(x^2 W_{BC}(x; u_1, \dots, u_r)^2 + 4\mu^2 \right)^{1/2}. \quad (23)$$

- F_4

$$W_{F_4}^{27} = \frac{a_1(x)}{24} - \frac{1}{2} \left\{ \left(-q + \sqrt{q^2 + 4p^3} \right)^{1/3} + \left(-q - \sqrt{q^2 + 4p^3} \right)^{1/3} \right\}, \quad (24)$$

where

$$\begin{aligned} p(x) &= -\frac{a_2}{6} - \frac{a_1^2}{144}, \\ q(x) &= \frac{1}{27} \left(\frac{a_1^3}{32} + \frac{9}{8} a_1 a_2 + 27 a_3 \right), \end{aligned} \quad (25)$$

and a_1, a_2 and a_3 are defined in (14).

- G_2

$$W_{G_2}^8 = \frac{1}{6} (p_1 + \sqrt{p_1^2 + 12p_2}), \quad (26)$$

where

$$p_1 = 6x^4 - 2ux^2, \quad p_2 = x^8 - 2ux^6 + u^2x^4 - vx^2 + 12\mu^4. \quad (27)$$

Note that for C_r and G_2 cases, the superpotentials $W_G^{\mathcal{R}}(x)$ depend on the scale parameter μ explicitly. So far we have seen the SW spectral curves for general gauge groups. The SW differential defined on these spectral curves take the simple form[3]:

$$\lambda_{SW} = \frac{1}{2\pi i} x \frac{dz}{z}. \quad (28)$$

In the next section we will study the period integrals of the SW differential using the two-dimensional topological LG theory.

3 Picard-Fuchs Equations and 2D Topological Landau-Ginzburg Models

In this section we consider the *ADE* gauge groups. To describe the moduli space of the Coulomb branch we adopt the flat coordinate system $(t_1, t_2, \dots, t_r)$ developed in the *A-D-E* singularity theory instead of the conventional Casimir coordinates $(u_1, u_2, \dots, u_r)$. The coordinate transformation is read off from the residue integral

$$t_i = c_i \oint dx W_G^{\mathcal{R}}(x, u)^{\frac{e_i}{h}}, \quad i = 1, \dots, r \quad (29)$$

with a suitable constant c_i [13, 15]. Notice that the overall degree of $W_G^{\mathcal{R}}$ is equal to h . The flat coordinates t_i are expressed as polynomials in u_i .

Firstly we discuss the role of flat coordinates in the two-dimensional topological Landau-Ginzburg models. We define the primary fields $\phi_i(x, t)$ as the derivatives of the superpotential $W_G(x, t)$:

$$\phi_i(x, t) = \frac{\partial W_G(x, t)}{\partial t_i}. \quad (30)$$

We choose the normalization factor c_r such that $\phi_r(x, t) = 1$. We now consider two-dimensional topological gravity coupled to topological LG model[16]. In this case, the primary field ϕ_r is regarded as the puncture operator P . In the flat coordinate system, the topological metric $\eta_{ij} = \langle \phi_i \phi_j P \rangle$ is independent of t_i and takes the form:

$$\eta_{ij} = \delta_{e_i + e_j, h}. \quad (31)$$

Furthermore, the primary fields obey the operator product expansions

$$\phi_i(x, t) \phi_j(x, t) = \sum_{k=1}^r C_{ij}^k(t) \phi_k(x, t) + Q_{ij}(x, t) \partial_x W_G(x, t). \quad (32)$$

The flatness condition implies that the function $Q_{ij}(x, t)$ in (32) satisfies

$$\frac{\partial^2 W_G(x, t)}{\partial t_i \partial t_j} = \partial_x Q_{ij}(x, t). \quad (33)$$

The structure constants $C_{ij}^k(t)$ are related to the three-point function $F_{ijk}(t) = \langle \phi_i \phi_j \phi_k \rangle$ by the relation $F_{ijk}(t) = \eta_{kl} C_{ij}^l(t)$. In two-dimensional topological theory, all the topological correlation functions are determined by the free energy $F(t)$. The three point function $F_{ijk}(t)$ is given by $\partial^3 F(t) / \partial t_i \partial t_j \partial t_k$.

The associativity of the chiral ring $(\phi_i \phi_j) \phi_k = \phi_i (\phi_j \phi_k)$ implies the relation $C_{ij}^l C_{lk}^n = C_{jk}^l C_{il}^n$. Let us introduce $r \times r$ matrices C_i , F_i and η by $(C_i)_j^k = C_{ij}^k$, $(F_i)_{jk} = F_{ijk}$ and $\eta = (\eta_{ij})$. The associativity condition leads to the commutativity of the matrices C_i ;

$$C_i C_j = C_j C_i. \quad (34)$$

Using $F_i = \eta C_i$, we obtain the Witten-Dijkgraaf-Verlinde-Verlinde (WDVV) equation:

$$F_i \eta^{-1} F_j = F_j \eta^{-1} F_i, \quad (35)$$

which is one of the important relations in two-dimensional topological theory.

Now we go back to the four-dimensional gauge theory. We consider the period integral of the SW differential λ_{SW} (28):

$$\Pi = \int_{\gamma} \lambda_{SW} \quad (36)$$

along the certain one-cycle γ on the spectral curve (16). In terms of the superpotential $W_G(x, t)$, the SW differential takes the form:

$$\lambda_{SW} = \frac{1}{2\pi i} \frac{x \partial_x W_G(x, t)}{\sqrt{W_G(x, t)^2 - 4\mu^2}} dx. \quad (37)$$

It is shown in [7] that Π obeys the set of differential equations:

$$\partial_{t_i} \partial_{t_j} \Pi = \sum_{k=1}^r C_{ij}^k(t) \partial_{t_k} \partial_{t_r} \Pi, \quad (38)$$

which is called as the Gauss-Manin system in the singularity theory.

In addition to the Gauss-Manin system, the SW differential satisfies another two differential equations. From the scaling relation to the superpotential

$$\left(\sum_{i=1}^r q_i t_i \partial_{t_i} + x \partial_x \right) W_G(x, t) = h W_G(x, t), \quad (39)$$

we obtain the scaling relation for the period Π :

$$\left(\sum_{i=1}^r q_i t_i \partial_{t_i} + h^\vee \mu \partial_\mu - 1 \right) \Pi = 0. \quad (40)$$

The final differential equation is obtained by regarding the l.h.s. of the spectral curve (16) as the superpotential of the topological CP^1 model [18]:

$$W_{CP^1}(z) = z + \frac{\mu^2}{z} - t_r. \quad (41)$$

Since $\log \mu^2$ and t_r are flat coordinates of the CP^1 model, we obtain the CP^1 relation:

$$\left((\mu \partial_\mu)^2 - 4\mu^2 \partial_{t_r}^2 \right) \Pi = 0. \quad (42)$$

Combining the scaling relation (40) and the CP^1 relation (42), we obtain the differential equation

$$\left\{ \left(\sum_{i=1}^r q_i t_i \frac{\partial}{\partial t_i} - 1 \right)^2 - 4\mu^2 h^2 \frac{\partial^2}{\partial t_r^2} \right\} \Pi = 0 \quad (43)$$

By solving the Gauss-Manin system (38) and the scaling equation (43), we may analyze the exact solutions in the Coulomb branch. For classical gauge groups, the present Picard-Fuchs equations are shown to be the same as those appeared in the previous works[19], in which various gauge theories with or without hypermultiplets are discussed.

In the weak coupling region where the scale parameter $\Lambda^{2h} = 4\mu^2$ are small, the solutions of these Picard-Fuchs equations are studied extensively using various methods[19, 20], which are shown to agree with the microscopic instanton calculation [21]. In the next section, we study the solutions in the strong coupling region. But before going to the next section, we discuss an important consequence of the Gauss-Manin system. Since the dual period a_{DI} also satisfies the Gauss-Manin system (38), this Gauss-Manin system provides the third-order differential equation for the prepotential $\mathcal{F}(a)$:

$$\tilde{\mathcal{F}}_{ijk} = \sum_{l=1}^r C_{ij}^l \tilde{\mathcal{F}}_{lrk}, \quad (44)$$

where

$$\begin{aligned} \tilde{\mathcal{F}}_{ijk} &= \partial_{t_i} a_I \partial_{t_j} a_J \partial_{t_k} a_K \mathcal{F}_{IJK}, \\ \mathcal{F}_{IJK} &= \partial_{a_I} \partial_{a_J} \partial_{a_K} \mathcal{F}(a) \end{aligned} \quad (45)$$

The equations (44) is very similar to $F_{ijk}(t) = C_{ij}^l \eta_{kl}$ in two-dimensional topological theory. Let us introduce matrices $\tilde{\mathcal{F}}_i$, $\mathcal{G}$ and $\mathcal{F}_I$ defined by $(\tilde{\mathcal{F}}_i)_{jk} = \tilde{\mathcal{F}}_{ijk}$, $\mathcal{G} = \tilde{\mathcal{F}}_r$, and $(\mathcal{F}_I)_{JK} = \mathcal{F}_{IJK}$, respectively. Then we find the WDVV equations in the Seiberg-Witten theory[22, 9]:

$$\tilde{\mathcal{F}}_i \mathcal{G}^{-1} \tilde{\mathcal{F}}_j = \tilde{\mathcal{F}}_j \mathcal{G}^{-1} \tilde{\mathcal{F}}_i, \quad (46)$$

which may be written in the form

$$\mathcal{F}_I \mathcal{F}_K^{-1} \mathcal{F}_J = \mathcal{F}_J \mathcal{F}_K^{-1} \mathcal{F}_I. \quad (47)$$

In addition to the WDVV equation, the prepotential satisfies the scaling equation[23]:

$$\left(\sum_{I=1}^r a_I \partial_{a_I} + h^\vee \mu \partial_\mu \right) \mathcal{F}(a) = 2\mathcal{F}(a). \quad (48)$$

This scaling equation is important to calculate the instanton correction to the prepotential in the weak coupling region[8]. Recently it is shown that the prepotential satisfies further non-trivial equations [24] obtained from the Whitham hierarchy. Instanton correction to the prepotential has been calculated in this framework[25] for some gauge theories. It would be interesting to study the relation between the WDVV equation approach and this formulation.

4 Superconformal point

One of interesting phenomena in the strong coupling physics of $N = 2$ supersymmetric gauge theory is the existence of non-trivial $N = 2$ superconformal fixed point in the Coulomb branch[11, 26, 27]. At this point, massless solitons of mutually nonlocal charges coexist. The superconformal field theory is characterized by the scaling operators with non-trivial (fractional) scaling dimensions. In particular, the calculations of the scaling exponents based on the exact solution suggest that the superconformal fixed points are characterized by the A - D - E classification[27]. We will study this RG fixed point in view of the Picard-Fuchs equations obtained in the previous section.

For $G = ADE$, the superconformal fixed points exist at

$$t_1 = \cdots = t_{r-1} = 0, \quad t_r = \pm 2\mu.$$

We take the plus sign without loss of generality. Let us introduce new flat coordinates $\tilde{t}_i$ by shifting t_r by 2μ :

$$t_i = \tilde{t}_i, \quad (i = 1, \cdots, r-1), \quad t_r = \tilde{t}_r + 2\mu \quad (49)$$

Since the OPE coefficients $C_{ij}^k(t)$ are independent of t_r [8], the Gauss-Manin system does not change its form under this change of coordinates:

$$\left(\partial_{\tilde{t}_i} \partial_{\tilde{t}_j} - \sum_{k=1}^r C_{ij}^k(\tilde{t}) \partial_{\tilde{t}_k} \partial_{\tilde{t}_r} \right) \Pi = 0, \quad (50)$$

The scaling equation (43), on the other hand, becomes

$$\left\{ \left(\sum_{i=1}^r q_i \tilde{t}_i \partial_{\tilde{t}_i} - 1 \right)^2 + 4\mu h \left[\left(\sum_{i=1}^r q_i \tilde{t}_i \partial_{\tilde{t}_i} - 1 \right) \partial_{\tilde{t}_r} + \frac{h}{2} \partial_{\tilde{t}_r} \right] \right\} \Pi = 0. \quad (51)$$

Introduce the scaling parameter ϵ by

$$\tilde{t}_i = \epsilon^{q_i} \rho_i, \quad (i = 1, \dots, r-1), \quad \tilde{t}_r = \epsilon^h \quad (52)$$

and consider the limit $\epsilon \rightarrow 0$. We are interested in the solution of the SW periods which behave like

$$\Pi = \epsilon^\alpha f(\rho) + \dots, \quad (53)$$

as $\epsilon \rightarrow 0$. Since

$$\partial_{\tilde{t}_i} = \epsilon^{-q_i} \partial_{\rho_i}, \quad \partial_{\tilde{t}_r} = \frac{1}{h} \epsilon^{-h} \left(\epsilon \partial_\epsilon - \sum_{i=1}^{r-1} q_i \rho_i \partial_{\rho_i} \right), \quad (54)$$

the Gauss-Manin system (50) for $i, j < r$ becomes the differential equations for $f(\rho)$ with respect to ρ :

$$\begin{aligned} & \left(\partial_{\rho_i} \partial_{\rho_j} - \frac{1}{h} \sum_{k=1}^{r-1} \tilde{C}_{ij}^k(\rho) \partial_{\rho_k} \left[\alpha - \sum_{l=1}^{r-1} q_l \rho_l \partial_{\rho_l} \right] \right. \\ & \left. - \frac{1}{h^2} \tilde{C}_{ij}^r(\rho) \left[\alpha - h - \sum_{l=1}^{r-1} q_l \rho_l \partial_{\rho_l} \right] \left[\alpha - \sum_{l=1}^{r-1} q_l \rho_l \partial_{\rho_l} \right] \right) f(\rho) = 0 \end{aligned} \quad (55)$$

where $\tilde{C}_{ij}^k(\rho) = C_{ij}^k(\tilde{t}) \epsilon^{q_i + q_j - q_k - h}$. As for the scaling equation (51), the second term is dominant in the superconformal limit. We thus find that $f(\rho)$ should satisfy the equation

$$\left(\alpha - \frac{h+2}{2} \right) \left[\alpha - \sum_{l=1}^{r-1} q_l \rho_l \partial_{\rho_l} \right] f(\rho) = 0 \quad (56)$$

This equation determines the exponent α in (53) such as

$$\alpha = \frac{h+2}{2}. \quad (57)$$

The superconformal field theory is characterized by the scaling operator $\text{tr} \phi^{q_i}$, whose conformal dimension is measured with respect to the BPS mass (1). From (53) and (57), we have

$$\langle \text{tr} \phi^{q_i} \rangle \sim (m_{BPS})^{\frac{2q_i}{h+2}} \quad (58)$$

Thus the scaling dimension of $\langle \text{tr} \phi^{q_i} \rangle$ is $2q_i/(h+2)$, which agrees with the result of [27].

We could examine the above arguments from the viewpoint of the SW curve. In terms of the parameters $\tilde{t}$, the SW curve becomes

$$z + \frac{\mu^2}{z} = W_G(x; t) = W_G(x; \tilde{t}) - 2\mu. \quad (59)$$

Let us introduce ξ by

$$\xi = \sqrt{z} + \frac{\mu}{\sqrt{z}}. \quad (60)$$

Then the curve is expressed in the form of the *ADE* superpotential with the Gaussian part ξ^2 :

$$\xi^2 = W_G(x; \tilde{t}). \quad (61)$$

The SW differential then becomes

$$\lambda_{SW} = \frac{1}{2\pi i} x \frac{dz}{z} = \frac{1}{2\pi i} 2x \frac{d\xi}{\sqrt{\xi^2 - 4\mu}}. \quad (62)$$

In the superconformal limit, expanding the above formula around $\epsilon = 0$ we obtain

$$\lambda_{SW} = -\frac{1}{2\pi i} \frac{1}{\sqrt{-\mu}} \sum_{n=0}^{\infty} \frac{(2n)!}{2^{2n}(n!)^2} \frac{W_G(x, \tilde{t})^{\frac{2n+1}{2}} dx}{(2n+1)(4\mu)^n}, \quad (63)$$

up to the total derivative term. After rescaling $x = \epsilon \tilde{x}$, the leading term in (63) is

$$\lambda_{SW} = -\frac{1}{2\pi i \sqrt{-\mu}} \epsilon^{\frac{h+2}{2}} \sqrt{W_G(\tilde{x}; \rho_1, \dots, \rho_{r-1}, 1)} d\tilde{x} + \dots, \quad (64)$$

which also leads to the exponent (57). Note that the derivative of the SW period integrals:

$$\frac{\partial \Pi}{\partial \rho_i} \sim \epsilon^{\frac{h+2}{2}} \int \frac{\partial_{\rho_i} W_G(\tilde{x}; \rho)}{\sqrt{W_G(\tilde{x}; \rho)}} d\tilde{x}$$

is the period of the curve

$$y^2 = W_G(\tilde{x}; \rho_1, \dots, \rho_{r-1}, 1). \quad (65)$$

Thus we find that the superconformal fixed point is simply characterized by the *ADE* superpotential $W_G(x, \tilde{t})$. The SW curve reduce to the hyperelliptic type curve (65). For example, let us consider the A_2 case. The SW curve (65) is given by

$$y^2 = x^3 - \rho x - 1,$$

which is nothing but the curve of $SU(2)$ gauge theory with $N_f = 1$ at the superconformal point (the small torus in [11]). The Gauss-Manin system (50) becomes

$$\left[(4\rho^3 - 27)\partial_\rho^2 - \frac{5}{4}\rho \right] f(\rho) = 0. \quad (66)$$

One may solve this equation around $\rho = 0$ found that the results agree with those obtained in [11].

5 Discussion

In this paper, we have seen the close relationship between the four-dimensional gauge theory with ADE gauge group and two-dimensional topological LG models coupled to topological gravity. We have examined the exact solutions around the superconformal points using the Picard-Fuchs equations and showed that the superconformal points are simply characterized by the $A-D-E$ singularity. For other gauge theories with or without matter hypermultiplets, on the other hand, it is difficult to extend the present results because of the complexity of the superpotential. But at the superconformal point we would expect that the curves become simple and are classified by the $A-D-E$ singularity[27]. The two-dimensional topological field theory would provide a useful tool for analyzing the physics in both weak and strong coupling region.

It seems interesting to compare the expansion (63) of the SW differential around the superconformal point with that around the origin ($t_i = 0$) of the Coulomb branch:

$$\lambda_{SW} = -\frac{1}{2\pi i} \frac{1}{\sqrt{-4\mu^2}} \sum_{n=0}^{\infty} \frac{(2n)!}{2^{2n}(n!)^2} \frac{W_G(x, t)^{2n+1} dx}{(2n+1)(4\mu^2)^n}. \quad (67)$$

In [10], it has been shown that the period integrals of (67) are expressed directly in terms of the one-point function $\langle \sigma_n(\phi_i) \rangle$ of the n -th gravitational descendant $\sigma_n(\phi_i)$ of the primary field ϕ_i in two-dimensional topological LG models coupled to topological gravity. The one-point function satisfies the same Gauss-Manin system, which is evaluated by the residue integrals[15]:

$$\langle \sigma_n(\phi_i) \rangle = b_{n,i} \sum_{j=1}^r \eta_{ij} \oint W_G(x, t)^{\frac{e_j}{h} + n + 1}, \quad (68)$$

where $b_{n,i}$ is certain constant. In this formulation, the Gauss-Manin system (38) is also derived from the topological recursion relation [17]:

$$\langle \sigma_n(\phi_i)XY \rangle = \sum_j \langle \sigma_{n-1}(\phi_i)\phi_j \rangle \langle \phi^j XY \rangle \quad (69)$$

and the puncture equation (in the small phase space):

$$\langle P \prod_{i=1}^s \sigma_{n_i}(\phi_{\alpha_i}) \rangle = \sum_{i=1}^s \langle \prod_{j=1}^s \sigma_{n_j - \delta_{ji}}(\phi_{\alpha_j}) \rangle. \quad (70)$$

One might expect that the similar relation would hold in the superconformal case, which would be important to understand the relation between the SW theory and $d < 1$ topological strings.

Acknowledgments The author would like to thank C.-S. Xiong and S.-K. Yang for useful discussions. This work is supported in part by the Grant-in-Aid from the Ministry of Education, Science and Culture, Priority Area: “Supersymmetry and Unified Theory of Elementary Particles” (#707).

References

- [1] N. Seiberg and E. Witten Nucl. Phys. **B426** (1994) 19, [hep-th/9407087](#); Nucl. Phys. **B431** (1994) 484, [hep-th/9408099](#)
- [2] A. Gorskii, I. Krichever, A. Marshakov, A. Mironov and A. Morozov, Phys. Lett. **B355** (1995) 466, [hep-th/9505035](#)
- [3] E.J. Martinec and N.P. Warner, Nucl. Phys. **B459** (1996) 97, [hep-th/9509161](#)
- [4] R. Donagi and E. Witten, Nucl. Phys. **B460** (1996) 299, [hep-th/9510101](#)
H. Itoyama and A. Morozov, Nucl. Phys. **B491** (1997) 529, [hep-th/9512161](#)
A. Gorsky, A. Marshakov, A. Mironov, A. Morozov, Phys. Lett. **B380** (1996) 75, [hep-th/9603140](#)
E. D'Hoker and D.H. Phong, Nucl. Phys. **B513** (1998) 405, [hep-th/9709053](#); Nucl. Phys. **B534** (1998) 697, [hep-th/9804126](#)
A.J. Bordner, E. Corrigan and R. Sasaki, Prog. Theor. Phys. **100** (1998) 1107, [hep-th/9805106](#)
A. J. Bordner, R. Sasaki and K. Takasaki, Prog. Theor. Phys. **101** (1999) 487, [hep-th/9809068](#)
A. Gorsky and A. Mironov, *Solutions to the reflection equation and integrable systems for $N = 2$ SQCD with classical groups*, [hep-th/9902030](#)
- [5] A. Klemm, W. Lerche, P. Mayr, C. Vafa and N. Warner, Nucl. Phys. **B477** (1996) 746, [hep-th/9604034](#)
For reviews, see W. Lerche, *Introduction to Seiberg-Witten Theory and its Stringy Origin*, [hep-th/9611190](#)
A. Klemm, *On the Geometry behind $N = 2$ Supersymmetric Effective Actions in Four Dimensions*, [hep-th/9705131](#)
- [6] E. Witten, Nucl. Phys. **B500** (1997) 3, [hep-th/9703066](#)
- [7] K. Ito and S.-K. Yang, Phys. Lett. **B415** (1997) 45, [hep-th/9708017](#)
- [8] K. Ito and S.-K. Yang, Int. J. Mod. Phys. **A13** (1998) 5373, [hep-th/9712018](#)

- [9] K. Ito and S.-K. Yang, Phys. Lett. **B433** (1998) 58, [hep-th/9803126](#)
- [10] K. Ito, C.-S. Xiong and S.-K. Yang, Phys. Lett. **B441** (1998) 155, [hep-th/9807183](#)
- [11] P. Argyres and M. Douglas, Nucl. Phys. **B448** (1995) 93, [hep-th/9505062](#)
- [12] W. Lerche and N.P. Warner, Phys. Lett. **B423** (1998) 79, [hep-th/9608183](#)
- [13] T. Eguchi and S.-K. Yang, Phys. Lett. **B394** (1997) 315, [hep-th/9612086](#)
- [14] J.B. Zuber, Mod. Phys. Lett. **A9** (1994) 749, [hep-th/9312209](#)
- [15] T. Eguchi, Y. Yamada and S.-K. Yang, Mod. Phys. Lett. **A8** (1993) 1627
- [16] R. Dijkgraaf, H. Verlinde and E. Verlinde, Nucl. Phys. **B352** (1991) 59
- [17] E. Witten, Nucl. Phys. **B340** (1989); Surv. Diff. Geom. **1** (1991) 243;
R. Dijkgraaf and E. Witten, Nucl. Phys. **B342** (1990) 486
- [18] B. Dubrovin, *Geometry of 2D Topological Field Theories*, [hep-th/9407018](#);
T. Eguchi, K. Hori and C.-S. Xiong, Int. J. Mod. Phys. **A12** (1997) 1743, [hep-th/9605225](#);
A. Givental, *Stationary Phase Integrals, Quantum Toda Lattices, Flag Manifolds and the Mirror Conjecture*, [alg-geom/9612001](#);
A. Takahashi, *Primitive Forms, Topological LG Models Coupled to Gravity and Mirror Symmetry*, [math.AG/9802059](#)
- [19] A. Klemm, W. Lerche and S. Theisen, Int. J. Mod. Phys. **A11** (1996) 1929, [hep-th/9505150](#)
K. Ito and S.-K. Yang, Phys. Lett. **B366** (1996) 165, [hep-th/9507144](#); in
“Frontiers In Quantum Field Theory” p. 331 (World Scientific, Singapore,1996), [hep-th/9603073](#)
M. Matone, Phys. Lett. **B357** (1995) 342, [hep-th/9506102](#)
Y. Ohta, J. Math. Phys. **37** (1996) 6074, [hep-th/9604051](#); **38** (1997) 682, [hep-th/9604059](#); **40** (1999) 1891, [hep-th/9809180](#); *Picard-Fuchs Ordinary Differential Systems in $N = 2$ Supersymmetric Yang-Mills Theories*, [hep-th/9812085](#)

- K. Ito and N. Sasakura, Nucl. Phys. **B484** (1997) 141, [hep-th/9608054](#)
- H. Ewen, K. Foerger and S. Theisen, Nucl. Phys. **B485** (1997) 63, [hep-th/9609062](#)
- H. Ewen and K. Foerger, Int. J. Mod. Phys. **A12** (1997) 4725, [hep-th/9610049](#);
- J.M. Isidro, A. Mukherjee, J.P. Nunes and H.J. Schnitzer, Nucl. Phys. **B492** (1997) 647, [hep-th/9609116](#); Int. J. Mod. Phys. **A13** (1998) 233, [hep-th/9703176](#); Nucl. Phys. **B502** (1997) 363, [hep-th/9704174](#);
- M. Alishahiha, Phys. Lett. **B398** (1997) 100, [hep-th/9609157](#); Phys. Lett. **B418** (1998) 317, [hep-th/9703186](#)
- K. Ito, Phys. Lett. **B406** (1997) 54, [hep-th/9703180](#)
- A.M. Ghezelbash and A. Shafiekhani, Mod. Phys. Lett. **A13** (1998) 527, [hep-th/9708073](#);
- A.M. Ghezelbash, Phys. Lett. **B423** (1998) 87, [hep-th/9710068](#)
- [20] E. D' Hoker, I.M. Krichever and D.H. Phong, Nucl. Phys. **B489** (1997) 179 [hep-th/9609041](#); **B489** (1997) 211 [hep-th/9609145](#)
- T. Masuda and H. Suzuki, Int. J. Mod. Phys. **A12** (1997) 3413, [hep-th/9609006](#);
- Int. J. Mod. Phys. **A13** (1998) 1495, [hep-th/9609065](#); Nucl. Phys. **B495** (1997) 149, [hep-th/9612240](#)
- T. Masuda, T. Sasaki and H. Suzuki, Int. J. Mod. Phys. **A13** (1998) 3121, [hep-th/9705166](#)
- [21] D. Finnell and P. Pouliot, Nucl. Phys. **B453** (1995) 225, [hep-th/9503115](#);
- H. Aoyama, T. Harano, M. Sato and S. Wada, Phys. Lett. **B388** (1996) 331, [hep-th/9607076](#)
- K. Ito and N. Sasakura, Phys. Lett. **B382** (1996) 95, [hep-th/9602073](#); Nucl. Phys. **B484** (1997) 141, [hep-th/9608054](#); Mod. Phys. Lett. **A12** (1997) 205, [hep-th/9609104](#)
- T. Harano and M. Sato, Nucl. Phys. **B484** (1997) 167, [hep-th/9608060](#)
- N. Dorey, V.V. Khoze and M.P. Mattis, Phys. Rev. **D54** (1996) 2921, [hep-th/9603136](#); 7832, [hep-th/9607202](#); Phys. Lett. **B388** (1996) 324, [hep-th/9607066](#); **B390** (1997) 205, [hep-th/9606199](#); **B396** (1997) 141, [hep-th/9611016](#); Nucl. Phys. **B492** (1997) 607;

- F. Fucito and G. Travaglini, Phys. Rev. **D55** (1997) 1099, [hep-th/9605215](#)
A. Yung, Nucl. Phys. **B485** (1997) 38, [hep-th/9605096](#)
M.J. Slater, Phys. Lett. **B403** (1997) 57, [hep-th/9701170](#)
- [22] A. Marshakov, A. Mironov and A. Morozov, Phys. Lett. **B389** (1996) 43, [hep-th/9607109](#); Mod. Phys. Lett. **A12** (1997) 773, [hep-th/9701014](#); *More Evidence for the WDVV Equations in $N = 2$ SUSY Yang-Mills Theories*, [hep-th/9701123](#);
G. Bonelli and M. Matone, Phys. Rev. Lett. **77** (1996) 4712, [hep-th/9605090](#);
G. Bertoldi and M. Matone, Phys. Lett. **B425** (1998) 104, [hep-th/9712039](#)
- [23] M. Matone, Phys. Lett. **B357** (1995) 342, [hep-th/9506102](#)
J. Sonnenschein, S. Theisen and S. Yankielowicz, Phys. Lett. **B367** (1996) 145, [hep-th/9510129](#)
T. Eguchi and S.-K. Yang, Mod. Phys. Lett. **A11** (1996) 131, [hep-th/9510183](#)
- [24] A. Gorskii, A. Marshakov, and A. Mironov, Nucl. Phys. **B527** (1998) 690, [hep-th/9802007](#)
- [25] J. D. Edelstein, M. Mariño and J. Mas, Nucl. Phys. **B541** (1999) 671, [hep-th/9805172](#)
J. D. Edelstein, M. Gómez-Reino and J. Mas, *Instanton corrections in $\mathcal{N} = 2$ supersymmetric theories with classical gauge groups and fundamental matter hypermultiplets*, [hep-th/9904087](#)
- [26] P. Argyres, R. Plesser, N. Seiberg and E. Witten, Nucl. Phys. **B461** (1996) 71, [hep-th/9511154](#)
- [27] T. Eguchi, K. Hori, K. Ito and S.-K. Yang, Nucl. Phys. **471** (1996) 430, [hep-th/9603002](#)
T. Eguchi and K. Hori, *$N=2$ Superconformal Field Theories in Four-dimensions and A-D-E Classification*, [hep-th/9607125](#)